

Automata and Formal Languages — Homework 11

Due 15.01.2019

Exercise 11.1

Let $\text{inf}(w)$ denote the set of letters occurring infinitely often in the infinite word w . Give Büchi automata and ω -regular expressions for the following ω -languages over $\Sigma = \{a, b, c\}$:

- (a) $L_1 = \{w \in \Sigma^\omega : \text{inf}(w) \subseteq \{a, b\}\}$,
- (b) $L_2 = \{w \in \Sigma^\omega : \text{inf}(w) = \{a, b\}\}$,
- (c) $L_3 = \{w \in \Sigma^\omega : \{a, b\} \subseteq \text{inf}(w)\}$,
- (d) $L_4 = \{w \in \Sigma^\omega : \text{inf}(w) = \{a, b, c\}\}$.

Exercise 11.2

Give *deterministic* Büchi automata recognizing the following ω -languages over $\Sigma = \{a, b, c\}$:

- (a) $L_1 = \{w \in \Sigma^\omega : w \text{ contains at least one } c\}$,
- (b) $L_2 = \{w \in \Sigma^\omega : \text{in } w, \text{ every } a \text{ is immediately followed by a } b\}$,
- (c) $L_3 = \{w \in \Sigma^\omega : \text{in } w, \text{ between two successive } a\text{'s there are at least two } b\text{'s}\}$.

Exercise 11.3

Give *deterministic* Rabin automata, Muller automata and parity automata for the following language:

$$L = \{w \in \{a, b\}^\omega : w \text{ contains finitely many } a\text{'s}\}.$$

Exercise 11.4

Prove or disprove:


- (a) For every Büchi automaton A , there exists a Büchi automaton B with a single initial state and such that $L_\omega(A) = L_\omega(B)$;
- (b) For every Büchi automaton A , there exists a Büchi automaton B with a single accepting state and such that $L_\omega(A) = L_\omega(B)$;
- (c) There exists a Büchi automaton recognizing the finite ω -language $\{w\}$ such that $w \in \{0, 1, \dots, 9\}^\omega$ and w_i is the i^{th} decimal of $\sqrt{2}$.

Exercise 11.5

Give a procedure that translates non-deterministic Rabin automata to non-deterministic Büchi automata.

Solution 11.1

(a) $(a + b + c)^*(a + b)^\omega$, and



★ It was asked in class whether there exists a deterministic Büchi automaton accepting L_1 . We show that it is *not* the case. For the sake of contradiction, suppose there exists a deterministic Büchi automaton $B = (Q, \Sigma, \delta, q_0, F)$ such that $L_\omega(B) = L_1$. Since $cb^\omega \in L_1$, B must visit F infinitely often when reading cb^ω . In particular, this implies the existence of $m_1 > 0$ and $q_1 \in F$ such that $q_0 \xrightarrow{cb^{m_1}} q_1$. Similarly, since $b^{m_1}cb^\omega \in L_1$, there exist $m_2 > 0$ and $q_2 \in F$ such that $q_0 \xrightarrow{cb^{m_1}cb^{m_2}} q_2$. Since B is deterministic, we have $q_0 \xrightarrow{cb^{m_1}} q_1 \xrightarrow{cb^{m_2}} q_2$. By repeating this argument $|Q|$ times, we can construct $m_1, m_2, \dots, m_{|Q|} > 0$ and $q_1, q_2, \dots, q_{|Q|} \in F$ such that

$$q_0 \xrightarrow{cb^{m_1}} q_1 \xrightarrow{cb^{m_2}} q_2 \cdots \xrightarrow{cb^{m_{|Q|}}} q_{|Q|}.$$


By the pigeonhole principle, there exist $0 \leq i < j \leq |Q|$ such that $q_i = q_j$. Let

$$u = cb^{m_1}cb^{m_2} \cdots cb^{m_i},$$


$$v = cb^{m_{i+1}}cb^{m_{i+2}} \cdots cb^{m_j}.$$

We have $q_0 \xrightarrow{u} q_i \xrightarrow{v} q_j \xrightarrow{v} q_i \xrightarrow{u} \cdots$ which implies that $uv^\omega \in L_\omega(B)$. This is a contradiction since $uv^\omega \notin L_1$. \square


(b) $(a + b + c)^*(aa^*bb^*)^\omega$, and




(c) $((b + c)^*a(a + c)^*b)^\omega$, and



or




(d) $((b+c)^*a(a+c)^*b(a+b)^*c)^\omega$, and




Solution 11.2


(a)




(b)



(c)




or simply,




Solution 11.3


- We give the following Rabin automaton with acceptance condition $\{(\{q_1\}, \{q_0\})\}$, i.e. where q_1 must be visited infinitely often and q_0 must be visited finitely often:



- We give the following Muller automaton with acceptance condition $\{\{q_1\}\}$, i.e. where precisely $\{q_1\}$ must be visited infinitely often:



- We give the following parity automaton with acceptance condition $(\{q_0\}, \{q_0, q_1\})$:



Solution 11.4

- (a) True. The construction for NFAs still work for Büchi automata.

Let $B = (Q, \Sigma, \delta, Q_0, F)$ be a Büchi automaton. We add a state to Q which acts as the single initial state. More formally, we define $B' = (Q \cup \{q_{\text{init}}\}, \Sigma, \delta', \{q_{\text{init}}\}, F)$ where

$$\delta'(q, a) = \begin{cases} \bigcup_{q_0 \in Q_0} \delta(q_0, a) & \text{if } q = q_{\text{init}}, \\ \delta(q, a) & \text{otherwise.} \end{cases}$$

We have $L_\omega(B) = L_\omega(B')$, since there exists $q_0 \in Q_0$ such that

$$q_0 \xrightarrow{a_1}_B q_1 \xrightarrow{a_2}_B q_2 \xrightarrow{a_3}_B \dots$$

if and only if

$$q_{\text{init}} \xrightarrow{a_1}_{B'} q_1 \xrightarrow{a_2}_{B'} q_2 \xrightarrow{a_3}_{B'} \dots$$

- (b) False. Let $L = \{a^\omega, b^\omega\}$. Suppose there exists a Büchi automaton $B = (Q, \{a, b\}, \delta, Q_0, F)$ such that $L_\omega(B) = L$ and $F = \{q\}$. Since $a^\omega \in L$, there exist $q_0 \in Q_0$, $m \geq 0$ and $n > 0$ such that

$$q_0 \xrightarrow{a^m} q \xrightarrow{a^n} q.$$

Similarly, since $b^\omega \in L$, there exist $q'_0 \in Q_0$, $m' \geq 0$ and $n' > 0$ such that

$$q'_0 \xrightarrow{b^{m'}} q \xrightarrow{b^{n'}} q.$$

This implies that

$$q_0 \xrightarrow{a^m} q \xrightarrow{b^{n'}} q \xrightarrow{b^{n'}} \dots$$

Therefore, $a^m(b^{n'})^\omega \in L$, which is a contradiction. □

- (c) False. Suppose there exists a Büchi automaton $B = (Q, \{0, 1, \dots, 9\}, \delta, Q_0, F)$ such that $L_\omega(B) = \{w\}$. There exist $u \in \{0, 1, \dots, 9\}^*$, $v \in \{0, 1, \dots, 9\}^+$, $q_0 \in Q_0$ and $q \in F$ such that

$$q_0 \xrightarrow{u} q \xrightarrow{v} q.$$

Therefore, $uv^\omega \in L_\omega(B)$ which implies that $w = uv^\omega$. Since w represents the decimals of π , we conclude that π is rational, which is a contradiction. \square