

# Syntax of propositional logic

An **atomic formula** has the form  $A_i$  where  $i = 1, 2, 3, \dots$

**Formulas** are defined by the following inductive process:

1. All atomic formulas are formulas
2. For every formula  $F$ ,  $\neg F$  is a formula.
3. For all formulas  $F$  und  $G$ ,  $(F \wedge G)$  and  $(F \vee G)$  are formulas.

For  $(F \wedge G)$  we say  $F$  and  $G$ , conjunction of  $F$  and  $G$


For  $(F \vee G)$  we say  $F$  or  $G$ , disjunction of  $F$  and  $G$

For  $\neg F$  we say not  $F$ , negation of  $F$

# Syntax tree of a formula


Every formula can be represented by a syntax tree.

Example:  $F = \neg((\neg A_4 \vee A_1) \wedge A_3)$



# Subformulas

The **subformulas** of a formula are the formulas corresponding to the subtrees of its syntax tree.



# Semantics of propositional logic (I)

The elements of the set  $\{0, 1\}$  are called **truth values**.

An **assignment** is a function  $\mathcal{A}: D \rightarrow \{0, 1\}$ , where  $D$  is any subset of the atomic formulas.

We extend  $\mathcal{A}$  to a function  $\hat{\mathcal{A}}: E \rightarrow \{0, 1\}$ , where  $E \supseteq D$  is the set of formulas that can be built up using only the atomic formulas from  $D$ .

# Semantics of propositional logic (II)

$$\begin{aligned}\hat{\mathcal{A}}(A) &= \mathcal{A}(A) \quad \text{if } A \in D \text{ is an atomic formula} \\ \hat{\mathcal{A}}((F \wedge G)) &= \begin{cases} 1 & \text{if } \hat{\mathcal{A}}(F) = 1 \text{ and } \hat{\mathcal{A}}(G) = 1 \\ 0 & \text{otherwise} \end{cases} \\ \hat{\mathcal{A}}((F \vee G)) &= \begin{cases} 1 & \text{if } \hat{\mathcal{A}}(F) = 1 \text{ or } \hat{\mathcal{A}}(G) = 1 \\ 0 & \text{otherwise} \end{cases} \\ \hat{\mathcal{A}}(\neg F) &= \begin{cases} 1 & \text{if } \hat{\mathcal{A}}(F) = 0 \\ 0 & \text{otherwise} \end{cases}\end{aligned}$$

We write  $\mathcal{A}$  instead of  $\hat{\mathcal{A}}$ .

# Truth tables (I)

We can compute  $\hat{A}$  with the help of [truth tables](#).

**Observe:**  $\hat{A}(F)$  depends only on the definition of  $\mathcal{A}$  on the atomic formulas that occur in  $F$ .

Tables for the operators  $\vee$ ,  $\wedge$ ,  $\neg$ :

$A$	$B$	$A$	$\vee$	$B$	$A$	$B$	$A$	$\wedge$	$B$	$A$	$\neg$	$A$
0	0	0	0	0	0	0	0	0	0	0	1	0
0	1	0	1	1	0	1	0	0	1	1	0	1
1	0	1	1	0	1	0	1	0	0			
1	1	1	1	1	1	1	1	1	1			

# Abbreviations

$A, B, C,$

$P, Q, R,$  or ... for  $A_1, A_2, A_3 \dots$

$(F_1 \rightarrow F_2)$  for  $(\neg F_1 \vee F_2)$

$(F_1 \leftrightarrow F_2)$  for  $((F_1 \wedge F_2) \vee (\neg F_1 \wedge \neg F_2))$

$(\bigvee_{i=1}^n F_i)$  for  $(\dots ((F_1 \vee F_2) \vee F_3) \vee \dots \vee F_n)$

$(\bigwedge_{i=1}^n F_i)$  for  $(\dots ((F_1 \wedge F_2) \wedge F_3) \wedge \dots \wedge F_n)$

$\top$  or **true** or 1 for  $(A_1 \vee \neg A_1)$

$\perp$  or **false** or 0 for  $(A_1 \wedge \neg A_1)$

# Truth tables (II)

Tables for the operators  $\rightarrow$ ,  $\leftrightarrow$ :

$A$	$B$	$A$	$\rightarrow$	$B$
0	0	0	1	0
0	1	0	1	1
1	0	1	0	0
1	1	1	1	1

$A$	$B$	$A$	$\leftrightarrow$	$B$
0	0	0	1	0
0	1	0	0	1
1	0	1	0	0
1	1	1	1	1

**Name:** *implication*

**Name:** *equivalence*

**Interpretation:** If  $A$  holds, then  $B$  holds.

**Interpretation:**  $A$  holds if and only if  $B$  holds.



# Beware!!!

$A \rightarrow B$  does **not** say, that  $A$  is a cause of  $B$ .

“Penguins swim  $\rightarrow$  Horses neigh”  
is true (in our world).

$A \rightarrow B$  does not say **anything** about the truth value of  $A$ .

“Ms. Merkel is a criminal  $\rightarrow$  Ms. Merkel should go to prison”  
is true (in our world).

A false statement implies **anything**.

“Penguins fly  $\rightarrow$  Mr. Obama is a criminal”  
is true (in our world).

# Formalizing natural language (I)

A device consists of two parts  $A$  and  $B$ , and a red light. We know that:

- $A$  or  $B$  (or both) are broken.
- If  $A$  is broken, then  $B$  is broken.
- If  $B$  is broken and the red light is on, then  $A$  is not broken.
- The red light is on.

We use the atomic formulas:  $RL$  (red light on),  $AB$  ( $A$  is broken),  $BB$  ( $B$  is broken), and formalize this situation by means of the formula

$$((AB \vee BB) \wedge (AB \rightarrow BB)) \wedge ((BB \wedge RO) \rightarrow \neg AB) \wedge RO$$

# Formalizing natural language (II)

Full truth table:

<i>RO</i>	<i>AB</i>	<i>BB</i>	$((AB \vee BB) \wedge (AB \rightarrow BB)) \wedge ((BB \wedge RO) \rightarrow \neg AB) \wedge RO$
0	0	0	0
0	0	1	0
0	1	0	0
0	1	1	0
1	0	0	0
1	0	1	1
1	1	0	0
1	1	1	0

# Formalizing natural language (III)

Formalize the Sudoku problem:

4				9				2
		1				5		
	9		3	4	5		1	
		8				2	5	
7		5		3		4	6	1
	4	6				9		8
	6		1	5	9		8	
		9				6		
5				7				4

An atomic formula  $X_{YZ}$  for each triple  $(X, Y, Z) \in \{1, \dots, 9\}^3$ :

$X_{YZ}$  = the square at row  $Y$  and column  $Z$  contains the number  $X$

**Example:** The first row contains all digits from 1 to 9

$$\bigwedge_{X=1}^9 \left( \bigvee_{Z=1}^9 X_{1Z} \right)$$

The truth table has

$2^{729}$  = 282401395870821749694910884220462786335135391185  
157752468340193086269383036119849990587392099522  
999697089786549828399657812329686587839094762655  
308848694610643079609148271612057263207249270352  
7723757359478834530365734912

rows.

# Models

Let  $F$  be a formula and let  $\mathcal{A}$  be an assignment.  $\mathcal{A}$  is **suitable** for  $F$  if it is defined for every atomic formula  $A_i$  occurring in  $F$ .

Let  $\mathcal{A}$  be suitable for  $F$ :

If  $\mathcal{A}(F) = 1$  then we write  $\mathcal{A} \models F$   
and say  $F$  holds under  $\mathcal{A}$   
or  $\mathcal{A}$  is a model of  $F$

If  $\mathcal{A}(F) = 0$  then we write  $\mathcal{A} \not\models F$   
and say  $F$  does not hold under  $\mathcal{A}$   
or  $\mathcal{A}$  is not a model of  $F$

# Validity and satisfiability

**Validity:** A formula  $F$  is **valid** (or a **tautology** if every suitable assignment for  $F$  is a model of  $F$ . We write  $\models F$  if  $F$  is valid, and  $\not\models F$  otherwise.

**Satisfiability:** A formula  $F$  is **satisfiable** if it has at least one model, otherwise  $F$  is **unsatisfiable**.

A (finite or infinite!) set of formulas  $S$  is **satisfiable** if there is an assignment that is a model of every formula in  $S$ .

# Exercise

	Valid	Satisfiable	Unsatisfiable
$A$			
$A \vee B$			
$A \vee \neg A$			
$A \wedge \neg A$			
$A \rightarrow \neg A$			
$A \rightarrow B$			
$A \rightarrow (B \rightarrow A)$			
$A \rightarrow (A \rightarrow B)$			
$A \leftrightarrow \neg A$			




# Exercise

Which of the following statements are true?

	J/N	C.ex.
If $F$ is valid, then $F$ is satisfiable		
If $F$ is satisfiable, then $\neg F$ is satisfiable		
If $F$ is valid, then $\neg F$ is satisfiable		
If $F$ is unsatisfiable, dann $\neg F$ is valid		

# Mirroring principle



# Consequence

A formula  $G$  is a **consequence** or **follows from** the formulas  $F_1, \dots, F_k$  if every model  $\mathcal{A}$  of  $F_1, \dots, F_k$  that is suitable for  $G$  is also a model of  $G$

If  $G$  is a consequence of  $F_1, \dots, F_k$  then we write  $F_1, \dots, F_k \models G$ .

# Consequence: example

$$\begin{aligned} & (AB \vee BB), (AB \rightarrow BB), \\ & ((BB \wedge RO) \rightarrow \neg AB), RO \models ((RO \wedge \neg AB) \wedge BB) \end{aligned}$$

# Exercise

$M$	$F$	$M \models F ?$
$A$	$A \vee B$	
$A$	$A \wedge B$	
$A, B$	$A \vee B$	
$A, B$	$A \wedge B$	
$A \wedge B$	$A$	
$A \vee B$	$A$	
$A, A \rightarrow B$	$B$	

# Consequence, validity, satisfiability

The following assertions are equivalent:

1.  $F_1, \dots, F_k \models G$ , e.g. ,  $G$  is a consequence of  $F_1, \dots, F_k$ .
2.  $((\bigwedge_{i=1}^k F_i) \rightarrow G)$  is valid.
3.  $((\bigwedge_{i=1}^k F_i) \wedge \neg G)$  is unsatisfiable.

# Exercise

Let  $S$  be a set of formulas, and let  $F$  and  $G$  be formulas. Which of the following assertions hold?

	Y/N
If $F$ satisfiable then $S \models F$ .	
If $F$ valid then $S \models F$ .	
If $F \in S$ then $S \models F$ .	
If $S \models F$ then $S \cup \{G\} \models F$ .	
$S \models F$ and $S \models \neg F$ cannot hold simultaneously.	
If $S \models G \rightarrow F$ and $S \models G$ then $S \models F$ .	