

We prove the undecidability of the validity problem for formulas of predicate logic with equality.

Recall: there is an algorithm that given a formula of predicate logic with equality returns a sat-equivalent formula of predicate logic.

It follows the validity problem for formulas of predicate logic with equality is also undecidable.

The proof is by reduction from the halting problem for **goto**-programs.

$Prog ::= \ell : Assign$	(assignment)
$\ell : \mathbf{goto} \ell'$	(unconditional jump)
$\ell : \mathbf{if} x_i \neq 0 \mathbf{then goto} \ell'$	(conditional jump)
$\ell : \mathbf{halt}$	(termination)
$Prog ; Prog$	(concatenation)

$Assign ::= x_i := 0 \mid x_i := x_j$
$x_i := x_j + 1 \mid x_i := x_j - 1$
$\ell ::= 1 \mid 2 \mid 3 \mid \dots$

Example

- 1: **if** $x_1 = 0$ **then goto** 4;
- 2: $x_1 := x_1 - 1$;
- 3: **goto** 1;
- 4: **halt**

Claim: **goto**-programs can simulate any program.

By the claim: a problem is decidable if it is solved by some **goto**-program.

We prove the following two theorems:

Theorem: The halting problem for **goto**-programs is undecidable: There is no (**goto**-)program that takes as input a **goto**-program P and a valuation β of the variables of P and decides whether P initialized with β terminates.

Theorem: If the validity problem is decidable, then the halting problem for **goto**-programs is decidable.

Fact: Programs and valuations can be encoded as integers.

Notations:

- $P(a_1, \dots, a_i)$ denotes the Program P initialized with $(a_1, \dots, a_i, 0, \dots, 0)$.
I.e., variables x_1, \dots, x_i are initialized with a_1, \dots, a_i and variables x_{i+1}, \dots, x_n with 0.
- Π_n denotes the program with code number n (if the program exists).

Fact: There exist computable encodings, i.e., encodings for which the following programs exist:

- Encoder.
Input: a program P .
Output: the code of P , i.e., the number n such that $P = \Pi_n$.
- Decoder.
Input: a number n .
Output: the program Π_n if n encodes a program, otherwise 'Not a program'.

The contradiction

Assumption: There is a program T such that for every pair $n, m \in \mathbb{N}$ the initialized program $T(n, m)$ halts and reports

Not a program	if n is not the code of a program
Yes	if n is the code of a program and $\Pi_n(m)$ halts
No	if n is the code of a program and $\Pi_n(m)$ does not halt

We show that this **assumption** leads to a contradiction.

Fact: The **assumption** implies the existence of a program T' such that for every $n \in \mathbb{N}$ the initialized program $T'(n)$

halts	if n is the code of a program and $\Pi_n(n)$ does not halt
does not halt	if n is not the code of a program or $\Pi_n(n)$ halts

Let k be the code of T' , i.e., $\Pi_k = T'$. Either the initialized program $T'(k)$ halts, or it does not halt. But:

- $T'(k)$ halts
- $\Rightarrow k$ is the code of a program and $\Pi_k(k)$ does not halt (Def. of T')
- $\Rightarrow T'(k)$ does not halt ($\Pi_k = T'$)
- $T'(k)$ does not halt
- $\Rightarrow \Pi_k(k)$ halts (Def. von T' , k is code)
- $\Rightarrow T'(k)$ halts ($\Pi_k = T'$)

So the **assumption is false**.

We assign to every program P and valuation β a formula $\phi_{P\beta}$ of predicate logic with equality such that

$\phi_{P\beta}$ is valid

if and only if

P with initialization β halts

There is a program that on input P, β outputs $\phi_{P\beta}$.

So no program can solve the validity problem.

Notations and definitions

Let k denote the number of instructions of P .
(The last instruction is always **halt**.)

Let n denote the number of variables of P .
(i.e., the variables of P are x_1, \dots, x_n .)

A **configuration** of P is a tuple $(pc, m_1, \dots, m_n) \in \mathbb{N}^{n+1}$.
 pc is the current value of the program counter and m_1, \dots, m_n the current valuation of the variables.

Convention: the successor of a configuration $(\ell_k, m_1, \dots, m_n)$ is again $(\ell_k, m_1, \dots, m_n)$.

Symbols of the formula $\phi_{P\beta}$

- R , predicate symbol of arity $(n + 2)$.
- $<$, predicate symbol of arity 2.
- f , function symbol of arity 1.
- 0 , constant.

- Universe: \mathbb{N} .
- $<^{\mathcal{A}}$ is the usual order on \mathbb{N} .
- $\mathbf{0}^{\mathcal{A}} = 0$.
- $f^{\mathcal{A}}$ is the successor function, i.e., $f^{\mathcal{A}}(n) = n + 1$.
- $R^{\mathcal{A}}(s, pc, m_1, \dots, m_n) = 1$ if (pc, m_1, \dots, m_n) is the configuration of P after s steps (for the initialization β).

$$\psi_{P\beta} = \psi_0 \wedge R(\mathbf{0}, \beta) \wedge \psi_1 \wedge \dots \wedge \psi_{k-1}$$

Meaning of $R(\mathbf{0}, \beta)$ in the structure \mathcal{A} : P is initialized with β

In the structure \mathcal{A} the formula ψ_i describes the effect of the i -th instruction of P . For instance:

- If i : $x_j := x_j + 1$ then

$$\psi_i = \forall x \forall y_1 \dots \forall y_n (\\ R(x, f^i(\mathbf{0}), y_1, \dots, y_n) \rightarrow \\ R(f(x), f^{(i+1)}(\mathbf{0}), y_1, \dots, y_{j-1}, f(y_j), y_{j+1}, \dots, y_n) \\)$$

1

1

- If i : **if $x_j = 0$ then goto j** then

$$\psi_i = \forall x \forall y_1 \dots \forall y_n (\\ R(x, f^i(\mathbf{0}), y_1, \dots, y_n) \rightarrow \\ (y_j = \mathbf{0} \wedge R(f(x), f^j(\mathbf{0}), y_1, \dots, y_n) \\ \vee \\ \neg(y_j = \mathbf{0}) \wedge R(f(x), f^{(i+1)}(\mathbf{0}), y_1, \dots, y_n) \\) \\)$$

ψ_0 guarantees that in every model the symbol $<$ is interpreted as a total order, that $\mathbf{0}$ is its smallest element, that $x < f(x)$ holds, and that $f(x)$ is the $<$ -successor of x :

$$\psi_0 = \forall x \forall y (x < y \wedge \neg(y < x)) \wedge \\ \forall x \forall y \forall z ((x < y \wedge y < z) \rightarrow x < z) \wedge \\ \forall x (\mathbf{0} < x \vee \mathbf{0} = x) \wedge \\ \forall x (x < f(x)) \wedge \\ \forall x \forall z (x < z \rightarrow (f(x) < z \vee f(x) = z))$$

1

1

The formula $\phi_{P\beta}$

We set

$$\phi_{P\beta} = \psi_{P\beta} \longrightarrow \exists x \exists y_1 \dots \exists y_n R(x, f^k(\mathbf{0}), y_1, \dots, y_n)$$

Theorem: $\phi_{P\beta}$ is valid iff program P with initialization β halts.

Proof: (\Rightarrow): If $\phi_{P\beta}$ is valid, then in particular the canonical structure \mathcal{A} is a model of $\phi_{P\beta}$. Since $\mathcal{A} \models \psi_{P\beta}$ clearly holds, we get $\mathcal{A} \models \exists x \exists y_1 \dots \exists y_n R(x, f^k(\mathbf{0}), y_1, \dots, y_n)$. So P initialized with β halts.

(\Leftarrow): (Sketch.) If $\phi_{P\beta}$ is not valid, then there is a structure $\mathcal{B} = (U_{\mathcal{B}}, I_{\mathcal{B}})$ such that

$$\mathcal{B} \models \psi_{P\beta} \text{ and } \mathcal{B} \not\models \exists x \exists y_1 \dots \exists y_n R(x, f^k(\mathbf{0}), y_1, \dots, y_n).$$

For every $i \geq 0$ let d_i be the element of $U_{\mathcal{B}}$ such that $(f^i(\mathbf{0}))^{\mathcal{B}} = d_i$. Since $\mathcal{B} \models \psi_{P\beta}$ we have $\mathcal{B} \models \psi_0$, and so (**why?**):

- $d_0 <^{\mathcal{B}} d_1 <^{\mathcal{B}} d_2 \dots$,
- $d_i = d_j$ iff $i = j$, and
- for every $d \in U_{\mathcal{B}}$: if $f^{\mathcal{B}}(d) = d_i$ then $d = d_{i-1}$.

Let (pc, m_1, \dots, m_n) be the configuration of P after s steps (with initialization β). Since $\mathcal{B} \models \psi_{P\beta}$ we have $R^{\mathcal{B}}(d^{s_i}, d^{z_i}, d^{m_{1i}}, \dots, d^{m_{ni}})$ for every $i \geq 0$. Since $\mathcal{B} \not\models \exists x \exists y_1 \dots \exists y_n R(x, f^k(\mathbf{0}), y_1, \dots, y_n)$, P does not terminate when initialized with β .

An alternative proof

The **tiling problem**:

Given: finite set of square tiles with fixed orientation and labelled borders: up, left, down, right. Each square is divided by its diagonals into four colored triangles.

Question: Can the plane be tiled with the given tiles in such a way, that neighbouring triangles in different tiles always have the same colour?

Theorem: The tiling problem is undecidable.

The reduction

We define for each set S of tiles a formula ϕ_S that is satisfiable iff the plane can be tiled with S .

Symbols: predicate symbol P_s of arity 2 for each tile $s \in S$,
function symbol f of arity 1.

Canonical structure \mathcal{A} :

- Universe: $\mathbb{Z} \times \mathbb{Z}$.
- $f^{\mathcal{A}}$ is the successor function, i.e., $f^{\mathcal{A}}(n) = n + 1$.
- $(i, j) \in P_s$ if tile s occupies the square with coordinates (i, j) .

Let H be the set of tile pairs (s, s') s.t. s' can be placed **right** from s .

Let V be the set of tile pairs (s, s') s.t. s' can be placed **above** s .

We take $\phi_S = \forall x \forall y (F_1 \wedge F_2)$ where

$$F_1 = \bigwedge_{s \neq s'} \neg (P_s(x, y) \wedge P_{s'}(x, y))$$

$$F_2 = \bigvee_{(s, s') \in H} (P_s(x, y) \wedge P_{s'}(f(x), y)) \wedge \bigvee_{(s, s') \in V} (P_s(x, y) \wedge P_{s'}(x, f(y)))$$

Corollary: The satisfiability problem is undecidable for closed formulas of the form $F = \forall x \forall y F^*$.

Corollary: The satisfiability problem is undecidable for closed formulas of the form $F = \forall x \exists z \forall y F^*$, where F^* contains no function symbols.

Prefix classes

We consider formulas in prenex form without function symbols.

Undecidable classes:

- $\forall^* \exists^*$ (Skolem, 1920)
- $\forall \forall \forall \exists$ (Suranyi, 1959)
- $\forall \exists \forall$ (Kahr, Moore, Wang, 1962)

Decidable classes:

- $\exists^* \forall^*$ (Bernays, Schönfinkel, 1928)
- $\exists^* \forall \exists^*$ (Ackerman, 1928)
- $\exists^* \forall^2 \exists^*$ (Gödel 1932, Kalmar 1933, Schütte 1934)